

## Anidulafungin

<b>Form:</b>	Vial containing 100mg dry powder
<b>Reconstitution:</b>	Reconstitute each vial with 30ml WFI and allow to stand for up to five minutes. <b>Further dilute before administration.</b>
<b>Administration Method:</b>	<p><b>IV Infusion</b></p> <p><b>Loading dose (Day 1 only):</b> Withdraw 50ml from 250ml infusion bag of compatible fluid and discard. Add 200mg (60ml) to remaining 200ml. Administer over 3 hours.</p> <p><b>Maintenance dose:</b> Add 100mg (30ml) to 100ml of compatible fluid. Administer over 90 minutes.</p> <p>Final concentration of 0.77mg/ml.</p>
<b>Compatibility &amp; Stability:</b>	Sodium Chloride 0.9% Glucose 5%
<b>Special Notes:</b>	<ul style="list-style-type: none"> <li>• Discard the solution if particulate matter or discoloration is present.</li> <li>• Infusion-related reactions have been reported with anidulafungin. Do not exceed the maximum infusion rate.</li> <li>• This is a RESTRICTED antimicrobial – refer to CUH Antimicrobial Guidelines or NCHD.ie app for further information.</li> </ul>

**Information provided relates to Ecalta<sup>®</sup> manufactured by Pfizer.**

*This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542*