

## Cyclimorph®

<b>Form:</b>	Cyclimorph 10® injection: Ampoules containing morphine tartrate 10mg and cyclizine tartrate 50mg in 1ml
<b>Reconstitution:</b>	Already in solution
<b>Administration Method:</b>	<p><b><u>IV Injection</u></b> Give by slow IV injection over a minimum of 4 - 5 minutes. The solution does not need to be diluted further. However, dilution with a small volume of diluent (5ml to 10ml) may aid administration.</p> <p><b><u>IM injection</u></b> No dilution required.</p> <p><b><u>SC Injection</u></b> No dilution required.</p>
<b>Compatibility &amp; Stability:</b>	Sodium Chloride 0.9% Glucose 5%
<b>Special Notes:</b>	<ul style="list-style-type: none"> <li>• Controlled Drug.</li> <li>• Monitor blood pressure and heart rate.</li> <li>• Monitor for respiratory depression.</li> <li>• Naloxone should be kept in all areas where opioids are administered.</li> <li>• Cyclimorph® should be administered as soon as possible after dilution. Immediately after dilution, and again just before injection, check the solution for signs of precipitation. Discard if there is any cloudiness or haze formation.</li> </ul>

**Information provided relates to Cyclimorph® manufactured by Amdipharm.**

*This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542*