

Labetalol

| | |
|---------------------------------------|--|
| Form: | 100mg per 20ml ampoule |
| Reconstitution: | Already in solution |
| Administration Method: | <p><u>IV Injection</u> Emergency use only. Use undiluted at a maximum rate of 50mg/min. Usual maximum total dose 200mg.</p> <p><u>IV infusion</u> Dilute to a concentration of 1mg/ml.</p> <p>Infuse the prescribed dosage using a rate-controlled infusion pump. Refer to CCU/Pharmacy for guidance.</p> |
| Compatibility & Stability: | Glucose 5% Sodium Chloride 0.9% & Glucose 5% |
| Special Notes: | <div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> <p style="text-align: center; color: blue;">For obstetric patients refer to CUMH guidelines or the Pharmacy Department</p> </div> <ul style="list-style-type: none"> • Monitor blood pressure and pulse regularly during infusion. |

Information provided relates to Trandate® manufactured by RPH Pharmaceuticals.

This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542