



Postural Hypotension



Cork Emergency Medicine

<http://EMed.ie>

What is postural hypotension?

Postural hypotension—or orthostatic hypotension— is when your blood pressure drops when you go from lying down to sitting up, or from sitting to standing.

When your blood pressure drops, less blood can go to your organs and muscles. This can make you more likely to fall.

What are the symptoms?

Although many people with postural hypotension have no symptoms, others do.

These symptoms can differ from person to person, and may include:

- Dizziness or lightheadedness
- Feeling about to faint, passing out, or falling
- Headaches, blurry or tunnel vision
- Feeling vague or muddled
- Feeling pressure across the back of your shoulders or neck
- Feeling nauseous, or hot and clammy
- Weakness or fatigue

When might it happen?

- When standing or sitting up suddenly
- In the morning when blood pressure is naturally lower
- After a large meal or alcohol
- During exercise
- When straining on the toilet
- When you are ill
- If you become anxious or panicky

If you are concerned, please contact the Emergency Department you first attended:

Mercy University Hospital
(021) 230 5011

M-UCC at SMHC (St. Mary's Health Campus) (021) 4926900

CUH
(021) 4920200

Local Injury Unit Mallow General Hospital
(022) 58506

Bantry General Hospital
(027) 52900



What causes postural hypotension?

Postural hypotension can be caused by or linked to:

- High blood pressure
- Diabetes, heart failure, atherosclerosis, or hardening of the arteries
- Taking some diuretics, antidepressants, or medicines to lower blood pressure
- Neurological conditions like Parkinson's disease and some types of dementia
- Dehydration
- Vitamin B12 deficiency or anemia
- Alcohol
- Prolonged bed rest

What can I do to prevent postural hypotension

- Tell your healthcare provider about any symptoms
- Ask if any of your medicines should be reduced or stopped
- Get out of bed slowly. First sit up, sit on the side of the bed, then stand up
- Take your time when changing position, such as when getting up from a chair
- Try to sit down when washing, showering, dressing, or working in the kitchen
- Exercise gently before getting up (move your feet up and down and clench and unclench your hands) or after standing (march in place)
- Make sure you have something to hold on to when you stand up
- Do not walk if you feel dizzy
- Drink 6-8 glasses of water or low-calorie drinks each day—unless you have been told to limit your fluid intake
- Avoid taking very hot baths or showers
- Try sleeping with extra pillows to raise your head

If you are concerned, please
contact the Emergency
Department you first attended:

**Mercy University
Hospital**
(021) 230 5011

**M-UCC at
SMHC (St.
Mary's
Health
Campus)**
(021)
4926900

CUH
(021) 4920200

**Local Injury Unit
Mallow General
Hospital**
(022) 58506

**Bantry General
Hospital**
(027) 52900

