

Early Management of Meningococcal Disease*

3RD EDITION - DECEMBER 2001

Estimate of child's weight (1-10 years)

Weight (kg) = 2 x (age in years + 4)

Systolic blood pressure = 80 + (age in years x 2)

N.B. Low BP is a pre-terminal sign in children

Conscious Level	Age	Normal Values Respiratory Rate	Heart Rate
Alert	<1	30-40	110-160
Responds to Voice	2-5	25-30	95-140
Responds to Pain	5-12	20-25	80-120
Unresponsive	>12	15-20	60-100

Observe HR, RR, BP, Perfusion, Conscious Level
Cardiac monitor and pulse oximetry. Take blood for Glucose, FBC, Clotting, U&E, Ca⁺⁺, Mg⁺⁺, PO₄, Blood cultures, Blood Gas (bicarb, base deficit), Cross-match

Colloid bolus (20ml/kg)

4.5% Human Albumin Solution (or Fresh Frozen Plasma or Hemacel/Gelofusine) i.v. or intra-osseous

Inotropes

Dopamine or Dobutamine at 10-20 mcg/kg/min (Make up 3 x weight (kg) mg in 50 ml 5% dextrose and run at 10 ml/hr = 10 mcg/kg/min) (these dilute solutions can be used via a peripheral vein)
Start Adrenaline via a central line only at 0.1 mcg/kg/min (Make up 300 mcg/kg in 50 ml of normal saline at 1 ml/hour=0.1 mcg/kg/min)

Intubation (call anaesthetist)

Atropine 20 mcg/kg (max 600 mcg) AND Thiopentone 3-5 mg/kg AND Suxamethonium 2 mg/kg (caution, high potassium) ETT size = age/4 + 4, ETT length (oral) = age/2 + 12, Then: morphine (100 mcg/kg) and midazolam (100 mcg/kg) every 30 mins

Hypoglycaemia (Glucose < 3 mmol/l)

5ml/kg 10% dextrose bolus i.v. and then dextrose infusion at 80% of maintenance requirements over 24 hours

Correction of metabolic acidosis pH < 7.2

1 mmol/kg NaHCO₃ i.v. = 1 ml/kg 8.4% NaHCO₃ over 20 mins or 2 ml/kg 4.2% NaHCO₃ in neonates

If K⁺ < 3.5 mmol/l

Give 0.25 mmol/kg over 30 mins i.v. with ECG monitoring
Caution if anuric

If total Calcium < 2 mmol/l or ionized Ca⁺⁺ < 1.0

Give 0.1 ml/kg 10% CaCl₂ (0.7 mmol/ml) over 30 mins i.v. (max 10 ml) or 0.3 ml/kg 10% Ca Gluconate (0.22 mmol/ml) over 30 mins (max 20 ml)

If Mg⁺⁺ < 0.75 mmol/l

Give 0.2 ml/kg of 50% MgSO₄ over 30 mins i.v. (max 10 ml)

Prophylaxis of household contacts

Inform Public Health Department, Give Rifampicin (bd for 2 days)
< 1yr 5 mg/kg • 1-12yrs 10 mg/kg • > 12yrs 600 mg or Ceftriaxone (single im dose)
< 12yrs 125 mg • > 12yrs 250 mg
or Ciprofloxacin as single 500 mg dose (adults only)

Diagnosis

Blood cultures, throat swab, whole blood (EDTA specimen) for PCR, rapid antigen test. Aspirations/scrapings from skin showing haemorrhagic rash

Serology

For suspected cases with no isolate or where PCR does not identify serogroup, clotted blood sample to MRU* (acute within 72 hrs and convalescent 10-28 days after presenting symptoms)

*PHLS Meningococcal Reference Unit

Tel: 0161 291 4628 Fax: 0161 446 2180 Out of hours: 0161 998 7070 (after 8pm)

© A.J. Pollard, S. Nadel, P. Habibi, S.N. Faust, I. Maconochie, N. Mehta, J. Britto, M. Levin (1998).
Department of Paediatrics, Imperial College School of Medicine, St Mary's Hospital, London W2
(*Arch Dis Child, March 1999; 80: 290-296)

Rev 12/01



RECOGNITION


May present with predominant SEPTICAEMIA (with shock), MENINGITIS (with raised ICP) or both. Purpuric/petechial non-blanching rash. Rash may be atypical or absent in some cases.

- Call consultant in A&E, Paediatrics, Anaesthesia or Intensive Care
- Initial assessment, looking for features of early shock/raised ICP
- DO NOT ATTEMPT LUMBAR PUNCTURE
- i.v. Cefotaxime (80 mg/kg) or Ceftriaxone (80 mg/kg)



Published by

Meningitis Research Foundation
Registered Charity No. 328205



www.meningitis.org