

Abatacept

Reduce direct handling to a minimum and wear appropriate personal protective equipment														
Abatacept dosing is weight based; ensure accuracy of documented weight before administration														
CAUTION: High Administration Risk Rating														
Form & Storage	Orencia® 250mg powder for concentrate for solution for infusion Pack includes a silicone free syringe	Refrigerate unopened vials at 2 - 8°C & protect from light.												
Reconstitution	<ul style="list-style-type: none">Using the silicone-free syringe provided, reconstitute each vial with 10mL water for injections, directing the stream to the wall of the vial.Remove the syringe and needle before swirling and rotating the vial gently to minimise foam formation; do not shake.Once the powder has dissolved, vent the vial with a needle to dissipate any foam. The reconstituted solution (25mg/mL) requires further dilution before administration.													
Compatibility & Stability	Sodium chloride 0.9%													
Administration	IV Infusion <ul style="list-style-type: none">Dilute required dose to a total volume of 100mL with sodium chloride 0.9%.Remove a volume of sodium chloride 0.9% from a 100mL infusion bag or bottle equal to the volume of the reconstituted dose required. <table><tr><th>Dose</th><th>Volume to remove from 100mL bag</th><th>Volume Orencia® to add to bag</th></tr><tr><td>500mg</td><td>20mL</td><td>20mL</td></tr><tr><td>750mg</td><td>30mL</td><td>30mL</td></tr><tr><td>1000mg</td><td>40mL</td><td>40mL</td></tr></table> <ul style="list-style-type: none">Using the same silicone-free disposable syringe as before, slowly add the reconstituted dose to the infusion container and gently mix the solution. The final concentration of abatacept should be no more than 10mg/mLGive over 30 minutes through a low-protein-binding filter (0.2 to 1.2micron).This filter B Braun Sterifix® 0.2µ Ref 4099303 is available to order from stores.		Dose	Volume to remove from 100mL bag	Volume Orencia® to add to bag	500mg	20mL	20mL	750mg	30mL	30mL	1000mg	40mL	40mL
Dose	Volume to remove from 100mL bag	Volume Orencia® to add to bag												
500mg	20mL	20mL												
750mg	30mL	30mL												
1000mg	40mL	40mL												
Documentation Requirements	Document batch numbers and expiry dates of vials in medical notes.													
Adverse Drug Reactions	Medicinal products for the treatment of hypersensitivity reactions, e.g. adrenaline, oxygen, antihistamines and corticosteroids should be available for immediate use in the event of an allergic reaction during administration of all infusions.													
Disposal	Dispose used vials, infusion bag and administration set in purple-lidded bins.													
Additional Information	<ul style="list-style-type: none">Orencia® contains maltose. Medicinal products containing maltose can interfere with the readings of blood glucose monitors that use test strips with glucose dehydrogenase pyrroloquinolinequinone (GDH-PQQ). ACCU-													

This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542

	<p>CHEK Inform II (stocked in CUH) that are labelled with a green symbol on the outer box do not have a clinically relevant maltose interference.</p> <ul style="list-style-type: none">• See PPG-CUH-CUH-243 <u>Policy Procedure and Guidelines for Management of Patients Attending CUH Infusion Unit for Intravenous Therapy CUH</u> for more information
--	---

Information relates to Orencia® (BMS)