

## Adrenaline (Epinephrine)

<b>CAUTION: High Administration Risk Rating</b>	
<b>Form</b>	1 in 10,000 (1mg per 10mL) prefilled syringe (Resuscitation trolley only)  1 in 1,000 (1mg per 1mL) ampoule
<b>Reconstitution</b>	<p>Prefilled syringe: Already in solution            Ampoule: Already in solution.</p> <ul style="list-style-type: none"> <li>• <b>Draw up using a 5 micron filter needle</b></li> <li>• <b>Use gloves when opening ampoules</b></li> </ul> <p><b><u>Dilute further before IV administration.</u></b></p>
<b>Compatibility &amp; Stability</b>	Sodium Chloride 0.9% Glucose 5%
<b>Administration</b>	<p><b><u>IV injection</u></b>            Use 1 in 10,000 (1mg per 10mL) prefilled syringe where available.</p> <p>If the prefilled syringe is not available the 1 in 1,000 (1mg per 1mL) preparation should be diluted to 1 in 10,000.            Dilute 1mL with 9mL Sodium Chloride 0.9% and mix well.</p> <p>Give by rapid IV injection. Administer via a central venous access device if already in place, or into a large peripheral vein.            IV injection administered via a peripheral vein should be followed by a 20mL flush of Sodium Chloride 0.9% to aid entry into the central circulation.</p> <p><b><u>IM Injection</u></b>            Use 1 in 1,000 (1mg per 1 mL) ampoule.</p> <p><b><u>IV infusion</u></b>            Contact ITU/Pharmacy for guidance.</p> <p>Discoloured solutions or solutions containing precipitate should not be used.</p>
<b>Extravasation</b>	Extravasation may cause tissue damage.
<b>Additional Information</b>	<ul style="list-style-type: none"> <li>• Repeated local administration may produce necrosis at the sites of injection.</li> <li>• Intramuscular injections of Adrenaline into the buttocks should be avoided because of the risk of tissue necrosis.</li> <li>• See <b>PPG-CUH-NUR-21</b> - Medication Protocol for the Administration of Epinephrine (Adrenaline) Injection BP 1:1000 by IM injection nurses and midwives for the management of a patient with anaphylaxis.</li> </ul>

**Information provided relates to Adrenaline manufactured by MercuryPharma and prefilled syringes manufactured by Aurum.**

*This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542*