

CefTAROLine fosamil

<p style="text-align: center;">SALAD</p> <p style="text-align: center;">Contains a PENICILLIN-like structure</p> <p style="text-align: center;">May be appropriate in penicillin-allergic patient. Refer to CUH Antimicrobial Guidelines on Eolas for further information before administration</p>	
<p style="text-align: center;">Restricted Antimicrobial</p> <p style="text-align: center;">See CUH Antimicrobial Guidelines on Eolas for further information</p>	
Form	Zinforo 600 mg powder for concentrate for solution for infusion
Reconstitution	Reconstitute each vial with 20mL WFI Shake well until solution is clear Dilute further before administration
Compatibility & Stability	Sodium chloride 0.9% Glucose 5%
Administration	<p>IV Infusion</p> <p>Add required dose to 100-250mL compatible infusion fluid</p> <p>Administer over 5 to 60 minutes for standard dose (every 12 hours) or 120 minutes for high dose (every 8 hours)</p> <p>The total time interval between starting reconstitution and completing preparation of the intravenous infusion should not exceed 30 minutes</p>
Monitoring	<p>Acute reactions</p> <ul style="list-style-type: none"> • anaphylaxis, hypersensitivity, • infusion site reactions (erythema, phlebitis, pain) • headache, dizziness • pyrexia • diarrhoea, nausea, vomiting, abdominal pain • rash, pruritis <p>Note: Contains arginine which may cause hypersensitivity reactions</p> <p>Monitor: infusion site, skin for urticaria, lip and face swelling, blood pressure, pulse, severe diarrhoea (colitis).</p>
Additional Information	<p>A 50ml infusion may be used if required (eg fluid restriction) but the residual volume in the infusion line must be flushed through at the same rate to avoid significant underdosing.</p> <p>Infusion related reactions can be managed by prolonging infusion duration.</p>

Information relates to Zinforo (Pfizer)