Information



Ceftolozane-Tazobactam

(Zerbaxa®)

SALAD Contains a PENICILLIN-like structure

May be appropriate	in penicillin-	allergic patient. Refer to CUH Antir her information before administrat	nicrobial Guidelines on Eolas for	
Please	e contact Mic	Restricted Antimicrobial robiology/ID/Antimicrobial pharma	cist for further information	
Form		Vial contains ceftolozane 1g and tazobactam 500 mg. Store vials at 2–8°C in fridge		
Reconstitution	1g cefto The fina	Add 10mL water for injections or sodium chloride 0.9% to each 1g ceftolozane/500mg tazobactam vial and shake gently. The final volume of each vial is approximately 11.4mL Dilute further prior to administration		
Compatibility & Stability		Sodium chloride 0.9% Glucose 5%		
Administration	•	IV infusion Any required dose to 100ml infusion fluid Administer over 60 minutes Dose of Volume of reconstituted injection		
		2g/1g	22.8ml (two vials)	
		1.5g/0.75g	17.1ml	
		1g/0.5g	11.4ml (one vial)	
Monitoring	Hyperse headach	Monitor: Blood pressure, heart rate. Hypersensitivity reactions including anaphylaxis, nausea, abdominal pain headache, dizziness, anxiety, fever, hypotension, tachycardia, rash, infusion site reactions, dyspnoea.		
Additional	Manufac	Manufacturer advises ceftolozane with tazobactam may influence driving and		

Information provided relates to Zerbaxa® manufactured by Merck Sharp & Dohme

performance of skilled tasks—increased risk of dizziness.