

Cyclizine

| | |
|--------------------------------------|--|
| Form | 50mg per 1mL ampoule |
| Reconstitution | <p>Already in solution</p> <ul style="list-style-type: none"> • Draw up using a 5 micron filter needle • Use gloves when opening ampoules |
| Compatibility & Stability | <p>Water for Injection Glucose 5% Sodium Chloride 0.9% - less stable</p> |
| Administration | <p>Immediately after dilution, and again just before injection, check the solution for signs of precipitation. Discard if there is any cloudiness or haze formation.</p> <p><u>IV Injection</u> Dilute solution with an equal volume of WFI and give slowly over at least 3 - 5 minutes.</p> <p><u>IM injection</u> No dilution required.</p> <p><u>Continuous SC Infusion(unlicensed)</u> Dilute with WFI only to required volume</p> |
| Extravasation | Extravasation is likely to cause tissue damage due to low pH. |
| Additional Information | <p>Administration via syringe driver is unlicensed and may increase the administration risk rating. To mitigate these risks:</p> <ul style="list-style-type: none"> • Contact the Pharmacy Department or Palliative care team for further guidance. • Consult the Palliative Care Formulary accessible on www.medicinescomplete.com or the Syringe Driver Survey Database (SDSD) (available after registration on www.palliattedrugs.com) for guidance on syringe driver compatibility. |

Information provided relates to Valoid® manufactured by Amdipharm.