

## Cyclizine

|                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                  |
|--------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|
| <b>Form</b>                          | 50mg per 1mL ampoule                                                                                                                                                                                                                                                                                                                                                                                                                                 | Store below 25°C |
| <b>Reconstitution</b>                | Already in solution <ul style="list-style-type: none"> <li>• Draw up using a 5 micron filter needle</li> <li>• Use gloves when opening ampoules</li> </ul>                                                                                                                                                                                                                                                                                           |                  |
| <b>Compatibility &amp; Stability</b> | Water for Injection<br>Glucose 5%<br>Sodium Chloride 0.9% - less stable                                                                                                                                                                                                                                                                                                                                                                              |                  |
| <b>Administration</b>                | <b>IV Injection</b>                                                                                                                                                                                                                                                                                                                                                                                                                                  |                  |
|                                      | Can be further diluted with 10mL of glucose 5% or sodium chloride 0.9% to aid slow administration (maximum concentration 5mg in 1mL).<br>Give slowly over 3 minutes<br>Immediately after dilution, and again just before injection, check the solution for signs of precipitation and cloudiness. Discard if there is any cloudiness or particulate matter present.                                                                                  |                  |
|                                      | <b>IM Injection</b>                                                                                                                                                                                                                                                                                                                                                                                                                                  |                  |
|                                      | No dilution required.                                                                                                                                                                                                                                                                                                                                                                                                                                |                  |
| <b>Monitoring</b>                    | <b>Continuous SC Infusion(unlicensed)</b>                                                                                                                                                                                                                                                                                                                                                                                                            |                  |
|                                      | Dilute with <b>WFI only</b> to required volume                                                                                                                                                                                                                                                                                                                                                                                                       |                  |
| <b>Monitoring</b>                    | Monitor BP, may cause hypotension<br>Injection site reactions including: vein tracking, erythema, pain, rash, thrombophlebitis and blisters.                                                                                                                                                                                                                                                                                                         |                  |
| <b>Extravasation</b>                 | Extravasation is likely to cause tissue damage due to low pH. If a central venous access device is unavailable, administer via a large peripheral vein monitoring insertion site closely using a recognised phlebitis scoring tool. Re-site cannula at first signs of inflammation                                                                                                                                                                   |                  |
| <b>Additional Information</b>        | Administration via syringe driver is unlicensed and may increase the administration risk rating. To mitigate these risks: <ul style="list-style-type: none"> <li>• Contact the Pharmacy Department or Palliative care team for further guidance.</li> <li>• Consult the <b>Palliative Care Formulary</b> or the <b>Drug Compatibility Checker</b> accessible via <a href="http://www.medicinescomplete.com">www.medicinescomplete.com</a></li> </ul> |                  |

**Information provided relates to Valoid® (Amdipharm) last updated 5/3/26**