

## Dalbavancin

<b>Restricted antimicrobial</b>									
Please contact Microbiology/ID/Antimicrobial pharmacist for further information									
<b>Form</b>	500mg per vial dry powder for concentrate for solution for infusion.								
<b>Reconstitution</b>	<ul style="list-style-type: none"> <li>Slowly add 25 mL water for injection to each vial</li> <li>Do not shake.</li> <li>To avoid foaming, alternate between gentle swirling and inversion of vial until contents dissolved completely (approx. 5 minutes).</li> </ul> <p style="text-align: center;"><b>Dilute further before administration</b></p>								
<b>Compatibility &amp; Stability</b>	Glucose 5% <b>ONLY</b>								
<b>Administration</b>	<p><b>IV Infusion</b></p> <p>Administer as an intravenous infusion over 30 minutes.</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr style="background-color: #e0f2f7;"> <th>Required Dose</th> <th>Volume of Glucose 5%</th> </tr> </thead> <tbody> <tr> <td>1500mg (3 vials)</td> <td>500mL</td> </tr> <tr> <td>1000mg (2 vials)</td> <td>250mL</td> </tr> <tr> <td>500mg (1 vial)</td> <td>100-250mL</td> </tr> </tbody> </table> <p>Infusion concentration should be between 1-5 mg/mL.</p>	Required Dose	Volume of Glucose 5%	1500mg (3 vials)	500mL	1000mg (2 vials)	250mL	500mg (1 vial)	100-250mL
Required Dose	Volume of Glucose 5%								
1500mg (3 vials)	500mL								
1000mg (2 vials)	250mL								
500mg (1 vial)	100-250mL								
<b>Monitoring</b>	Rapid administration can cause reactions including flushing of the upper body, urticaria, pruritis and/or rash. Stopping or slowing the infusion may result in cessation of these reactions.								
<b>Extravasation</b>	Dalbavancin has a low pH and may cause venous irritation and tissue damage in cases of extravasation. If a central venous access device is unavailable, administer via a large peripheral vein monitoring insertion site closely using a recognised phlebitis scoring tool. Re-site cannula at first signs of inflammation.								
<b>Additional Information</b>	<ul style="list-style-type: none"> <li>If a common intravenous line is being used to administer other drugs in addition to dalbavancin, the line should be flushed before and after each dalbavancin administration with glucose 5% solution for infusion.</li> <li>Do not mix dalbavancin with any other medicinal products or intravenous solutions.</li> </ul>								

**Information provided relates to Dalbavancin (Xydalba®) manufactured by AbbVie.**