

## CeFUroxime

<p><b>SALAD</b></p> <p><b>Contains a PENICILLIN-like structure</b></p> <p>May be appropriate in penicillin-allergic patient. Refer to CUH Antimicrobial Guidelines on <a href="#">Microguide</a> for further information before administration</p>	
<b>Form</b>	250mg, 750mg and 1.5 g dry powder vials
<b>Reconstitution</b>	<p><b><u>Intravenous</u></b> Add at least 2mL WFI to 250mg vial. Add at least 6mL WFI to 750mg vial. Add at least 15mL WFI to 1.5g vial.</p> <p><b><u>Intramuscular</u></b> Add 1mL WFI to 250mg vial. Add 3mL WFI to 750mg vial.</p>
<b>Compatibility &amp; Stability</b>	<p>Sodium Chloride 0.9% Glucose 5%</p> <p><b>From a microbiological point of view, should be used immediately;</b> however:</p> <ul style="list-style-type: none"> <li>• Reconstituted vials may be stored at 2–8°C for 24 hours.</li> <li>• Prepared infusions may be stored at 2–8°C and infused (at room temperature) within 24 hours.</li> </ul>
<b>Administration</b>	<p><b><u>IV Injection</u></b> Give slowly over 3 - 5 minutes.</p> <p><b><u>IV Infusion</u></b> After reconstitution, dilute required dose in 50 - 100mL of compatible fluid. Infuse over 30 - 60 minutes.</p> <p><b><u>IM injection</u></b> Not more than 750 mg should be injected at one site. For doses greater than 1.5 g intravenous administration should be used.</p>

**Information provided relates to Cefuroxime manufactured by Fresenius Kabi and GlaxoSmithKline.**