

## Cyclizine

<b>Form</b>	50mg per 1mL ampoule
<b>Reconstitution</b>	Already in solution
<b>Compatibility &amp; Stability</b>	Water for Injection Glucose 5% Sodium Chloride 0.9% - less stable
<b>Administration</b>	<p>Immediately after dilution, and again just before injection, check the solution for signs of precipitation. Discard if there is any cloudiness or haze formation.</p> <p><b><u>IV Injection</u></b> Dilute solution with an equal volume of WFI and give slowly over at least 3 - 5 minutes.</p> <p><b><u>IM injection</u></b> No dilution required.</p> <p><b><u>Continuous SC Infusion(unlicensed)</u></b> Dilute with <b>WFI only</b> to required volume</p>
<b>Extravasation</b>	Extravasation is likely to cause tissue damage due to low pH.
<b>Additional Information</b>	<p>Administration via syringe driver is unlicensed and may increase the administration risk rating. To mitigate these risks:</p> <ul style="list-style-type: none"> <li>• Contact the Pharmacy Department or Palliative care team for further guidance.</li> <li>• Consult the Palliative Care Formulary accessible on <a href="http://www.medicinescomplete.com">www.medicinescomplete.com</a> or the Syringe Driver Survey Database (<a href="http://www.palliativedrugs.com">SDSD</a>) (available after registration on <a href="http://www.palliativedrugs.com">www.palliativedrugs.com</a>) for <a href="#">guidance on syringe driver compatibility</a>.</li> </ul>

**Information provided relates to Valoid® manufactured by Amdipharm.**