

## Hydrocortisone (Solu-Cortef®)

<b>Form</b>	100mg dry powder vial as Hydrocortisone Sodium Succinate
<b>Reconstitution</b>	<p>Add 2mL WFI to each 100mg vial.</p> <p>Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Use solution only if it is clear. Reconstituted solution should be used immediately.</p>
<b>Compatibility &amp; Stability</b>	<p>Sodium Chloride 0.9%</p> <p>Glucose 5%</p>
<b>Administration Method</b>	<p><b><u>IV Injection</u></b> Give over 1 - 10 minutes.</p> <p><b><u>IV Infusion</u></b> Add reconstituted solution to at least 100mL of compatible fluid. Give over 20 - 30 minutes.</p> <p><b><u>IM Injection</u></b> No further dilution of reconstituted solution required.</p>
<b>Monitoring</b>	Monitor serum Na, K, Ca.
<b>Additional Information</b>	<ul style="list-style-type: none"> <li>Central serous chorioretinopathy is a retinal disorder that has been linked to the systemic use of corticosteroids. Patients should be advised to report any blurred vision or other visual disturbances with corticosteroid treatment.</li> </ul>

**Information provided relates to Solu-Cortef® manufactured by Pfizer.**