

Lidocaine

Potential SALAD Check strength . Also available as Lidocaine 1%																															
CAUTION: High Administration Risk Rating																															
Form	Lidocaine 2% (100mg per 5 mL) ampoules																														
Reconstitution	Already in solution																														
Compatibility & Stability	Glucose 5% Sodium Chloride 0.9%																														
Administration	<p>IV Injection Give 50 - 100mg over 2 minutes and flush immediately with 20mL sodium chloride 0.9%.</p> <p>IV Infusion Infusions of 2mg/mL generally used, but up to 8mg/mL if fluid restricted. Preferably administer via a central venous access device to avoid potential venous irritation. If given peripherally, choose a large vein and monitor the injection site closely.</p> <ul style="list-style-type: none"> For 2mg/mL solution (1g in 500mL) Add 50mL of 2% Lidocaine to 450mL of compatible infusion fluid to give 500mL of a solution containing 2mg/mL Lidocaine. <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Dose mg/min</th> <th>Rate mL/hour</th> </tr> </thead> <tbody> <tr><td>1</td><td>30</td></tr> <tr><td>2</td><td>60</td></tr> <tr><td>3</td><td>90</td></tr> <tr><td>4</td><td>120</td></tr> </tbody> </table> For 4mg/ml solution (2g in 500mL) Add 100mL of 2% Lidocaine to 400mL of compatible infusion fluid to give 500mL of a solution containing 4mg/mL Lidocaine. <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Dose mg/min</th> <th>Rate mL/hour</th> </tr> </thead> <tbody> <tr><td>1</td><td>15</td></tr> <tr><td>2</td><td>30</td></tr> <tr><td>3</td><td>45</td></tr> <tr><td>4</td><td>60</td></tr> </tbody> </table> For 8mg/ml solution (400mg in 50mL) Add 20mL of 2% Lidocaine to 30mL of compatible infusion fluid to give 50mL of a solution containing 8mg/mL Lidocaine. This may be used with a syringe pump in fluid restricted patients. <table border="1" style="margin-left: 20px;"> <thead> <tr> <th>Dose mg/min</th> <th>Rate mL/hour</th> </tr> </thead> <tbody> <tr><td>1</td><td>7.5</td></tr> <tr><td>2</td><td>15</td></tr> <tr><td>3</td><td>22.5</td></tr> <tr><td>4</td><td>30</td></tr> </tbody> </table> 	Dose mg/min	Rate mL/hour	1	30	2	60	3	90	4	120	Dose mg/min	Rate mL/hour	1	15	2	30	3	45	4	60	Dose mg/min	Rate mL/hour	1	7.5	2	15	3	22.5	4	30
Dose mg/min	Rate mL/hour																														
1	30																														
2	60																														
3	90																														
4	120																														
Dose mg/min	Rate mL/hour																														
1	15																														
2	30																														
3	45																														
4	60																														
Dose mg/min	Rate mL/hour																														
1	7.5																														
2	15																														
3	22.5																														
4	30																														
Monitoring	ECG monitoring is required.																														
Extravasation	Extravasation is likely to cause tissue damage due to acidic pH (<5).																														
Additional Information	Lidocaine products containing adrenaline or preservatives must not be given by IV injection.																														

Information provided relates to Lidocaine Mini-Plasco® manufactured by B Braun.

This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542