

Tobramycin

Form:	80mg per 2ml vial	
Reconstitution:	Already in Solution	
Administration Method:	Multiple Daily Dosing	Once Daily Dosing
	<u>IV Infusion</u> Dilute in 50 - 100ml compatible fluid and give over 20 - 60 minutes.	<u>IV Infusion</u> Dilute to 100ml compatible fluid and give over 60 minutes.
	<u>IV Injection</u> Slow Injection over 3 - 5 minutes.	<u>IV Injection</u> Not recommended
Extravasation	Extravasation may cause tissue damage due to its low pH.	
Compatibility & Stability:	Sodium Chloride 0.9% Glucose 5%	
Special Notes:	<ul style="list-style-type: none"> • Use Ideal Body Weight to calculate dose in obese patients – see CUH antimicrobial guidelines or NCHD.ie app for further information. • Plasma level monitoring recommended; refer to CUH antimicrobial guidelines or NCHD.ie app for further information. • This is a RESTRICTED antimicrobial- see CUH antimicrobial guidelines or NCHD.ie app for further information. 	

Information provided relates to Tobramycin manufactured by Hospira.

This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542