

## Ibuprofen

<b>SALAD</b>		
Caution with other BBraun products; Metronidazole 500mg in 100mL bottle		
<b>Approved for use in Theatres ONLY in CUH</b>		
<b>Form</b>	400mg solution for infusion in 100mL bottle	Store at room temperature in outer box for light protection.
<b>Reconstitution</b>	Already in solution – ready to administer	
<b>Compatibility &amp; Stability</b>	N/A	
<b>Administration</b>	<b>IV Infusion</b>	
	Administer over 30 minutes	
<b>Monitoring</b>	<ul style="list-style-type: none"> <li>• Monitor for signs of gastrointestinal bleeding, ulceration or perforation</li> <li>• Monitor for signs of bronchospasm, urticaria or angioedema</li> <li>• Monitor for oedema, hypertension and cardiac failure</li> <li>• infusion site reactions: pain and burning sensation, swelling, haematoma</li> </ul>	
<b>Additional Information</b>	The licensed maximum IV treatment duration is 3 days. Switch to oral treatment as soon as possible. Maintain adequate hydration to minimise risk of renal adverse effects	

### Information provided relates to Ibuprofen (BBraun)

*This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542*