

Pabrinex® (Vitamins B & C)

Form	<p>Vitamin B and C (paired ampoules) 2 x 5ml</p> <p>Each No. 1 ampoule (5mL) contains: Thiamine Hydrochloride 250mg Riboflavin (as Phosphate Sodium) 4mg Pyridoxine Hydrochloride 50mg</p> <p>Each No. 2 ampoule (5mL) contains: Ascorbic acid 500mg Nicotinamide 160mg Glucose (as monohydrate) 1000mg</p>
Reconstitution	<p>Already in solution</p> <ul style="list-style-type: none"> • Draw up using a 5micron filter needle • Use gloves when opening ampoules <p>Dilute further before administration.</p>
Compatibility & Stability	<p>Sodium Chloride 0.9%</p> <p>Glucose 5%</p>
Administration	<p><u>Intermittent IV infusion</u></p> <p>Draw up contents of two ampoules/one pair (1&2) into the same syringe, mix and add to 100mL infusion fluid.</p> <p>Infuse over at least 30 minutes.</p> <p>Up to three pairs of ampoules may be added to one bag. (One pair = Ampoule 1 + Ampoule 2) Administer immediately after the addition of ampoules to infusion fluid.</p>
Additional Information	<p>Risk of anaphylaxis is greatly reduced if Pabrinex® is given over at least 30 minutes. Facilities for treating anaphylaxis should be available.</p>

Information provided relates to Pabrinex® manufactured by Archimedes Pharmaceuticals.