

Solvito N[®]

Form	Dry powder vial Solvito N [®] contains thiamine, riboflavin, nicotinamide, pyridoxine, pantothenic acid, biotin, folic acid, cyanocobalamin, vitamin C.
Reconstitution	Dissolve with 10mL of water for injection and shake vigorously Dilute further before administration.
Compatibility & Stability	Glucose 5% (See notes below for compatibility with sodium chloride 0.9%)
Administration Method	<u>Peripheral or central IV route</u> Add reconstituted solution to 100mL Glucose 5% and infuse over a minimum period of 2-3 hours.
Additional Information	<div style="border: 1px solid blue; padding: 5px; margin-bottom: 10px;"> <p style="color: blue; text-align: center;">For obstetric patients refer to CUMH guidelines or the Pharmacy Department</p> </div> <ul style="list-style-type: none"> • Solvito N[®] is normally administered with Parenteral Nutrition. • For patients prescribed Additrace[®], Solivito N[®], and Vitlipid N Adult[®], or a combination of these, they can be infused together in 100mL glucose 5% or sodium chloride 0.9% over 2-3 hours.

Information provided relates to Solvito N[®] manufactured by Fresenius Kabi.

This information has been summarised to act as a guide for those administering IV medication. The monograph should be used in conjunction with the drug data sheet and BNF for information on dose, adverse effects, cautions and contra-indications. Further information is available from Pharmacy on 22146 or 22542